

CODIMENSION AND PROJECTIVE DIMENSION UP TO SYMMETRY

DINH VAN LE, UWE NAGEL, HOP D. NGUYEN, AND TIM RÖMER

ABSTRACT. Symmetric ideals in increasingly larger polynomial rings that form an ascending chain are investigated. We focus on the asymptotic behavior of codimensions and projective dimensions of ideals in such a chain. If the ideals are graded it is known that the codimensions grow eventually linearly. Here this result is extended to chains of arbitrary symmetric ideals. Moreover, the slope of the linear function is explicitly determined. We conjecture that the projective dimensions also grow eventually linearly. As part of the evidence we establish two non-trivial lower linear bounds of the projective dimensions for chains of monomial ideals. As an application, this yields Cohen-Macaulayness obstructions.

1. INTRODUCTION

Ascending chains of ideals that are invariant under actions of symmetric groups have recently attracted considerable attention. They arise naturally in various areas of mathematics, such as algebraic chemistry [1, 10], group theory [8], representation theory [7, 23, 24, 25, 26], toric algebra and algebraic statistics [2, 11, 12, 13, 16, 17, 18, 27], which provide frameworks and motivations for further studies. In [21] we investigated the behavior of the Castelnuovo-Mumford regularity along graded ideals in such a chain. Here we study the analogous problem for codimension and projective dimension.

Let \mathbb{N} denote the set of positive integers. Throughout the paper, fix an integer $c \in \mathbb{N}$ and any field K . For each $n \in \mathbb{N}$, let

$$R_n = K[x_{k,j} \mid 1 \leq k \leq c, 1 \leq j \leq n]$$

be the polynomial ring in $c \times n$ variables over K . These form an ascending chain

$$R_1 \subseteq R_2 \subseteq \cdots \subseteq R_n \subseteq \cdots$$

Let $\text{Sym}(n)$ denote the symmetric group on $\{1, \dots, n\}$. Considering it as stabilizer of $n+1$ in $\text{Sym}(n+1)$, similarly one gets an ascending chain of symmetric groups. Define an action of $\text{Sym}(n)$ on R_n induced by

$$\sigma \cdot x_{k,j} = x_{k,\sigma(j)} \quad \text{for every } \sigma \in \text{Sym}(n), 1 \leq k \leq c, 1 \leq j \leq n.$$

A sequence of ideals $(I_n)_{n \geq 1}$ with $I_n \subseteq R_n$ is called *Sym-invariant* if

$$\text{Sym}(n)(I_m) \subseteq I_n \quad \text{for all } m \leq n.$$

Observe that these ideals form an ascending chain as $I_n \cdot R_{n+1} \subseteq I_{n+1}$.

2010 *Mathematics Subject Classification.* 13A50, 13C15, 13D02, 13F20, 16P70, 16W22.

Key words and phrases. Invariant ideal, monoid, polynomial ring, symmetric group.

The second author was partially supported by Simons Foundation grant #317096. The third author was partially supported by a postdoctoral fellowship from the Vietnam Institute for Advanced Study in Mathematics (VIASM).

Even if one is primarily interested in Sym-invariant chains it is more convenient to work with a larger class of invariant objects, namely Inc^i -invariant chains, where Inc^i denotes a certain monoid of increasing functions on \mathbb{N} (see Section 2 for more details).

In [22], the second and fourth author introduced Hilbert series for Inc^i -invariant chains and proved that these series are rational (see also [20, Theorem 4.3] for another approach and [15, Theorems 2.4 and 3.3] for some explicit results in a special case). As a consequence, they determined the asymptotic behavior of the Krull dimension and multiplicity of graded ideals in an Inc^i -invariant chain: the Krull dimension grows eventually linearly, whereas the multiplicity grows eventually exponentially. This result motivates a more general line of investigations:

Problem 1.1. *Study the asymptotic behavior of invariants of ideals in Sym-invariant or, more generally, Inc^i -invariant chains.*

In [21], this problem was studied in the case of the Castelnuovo-Mumford regularity. There we conjectured that this invariant grows eventually linearly and provided some evidence supporting this conjecture. In particular, a linear upper bound for the Castelnuovo-Mumford regularity of graded ideals was established. As mentioned above, the present work studies the asymptotic behavior of codimensions (i.e. heights) and projective dimensions of ideals in Inc^i -invariant chains.

For an Inc^i -invariant chain of graded ideals $(I_n)_{n \geq 1}$, it follows from [22, Theorem 7.10] that $\text{codim} I_n$ is eventually a linear function. However, not much is known about this function. Here, we extend this result to Inc^i -invariant chains of ideals that are not necessarily graded. More importantly, our new approach also produces an explicit description for the leading coefficient of the linear function (see Theorem 3.8).

To motivate our study on the asymptotic behavior of the projective dimension, let us consider a simple example.

Example 1.2. Let $(I_n)_{n \geq 1}$ be an Inc^1 -invariant chain with

$$I_n = \begin{cases} \langle 0 \rangle & \text{if } n = 1, 2, 3, \\ \langle x_{1,2}^3, x_{1,4}^2 x_{2,1}, x_{2,2} x_{3,3} \rangle & \text{if } n = 4, \\ \langle \text{Inc}_{4,n}^1(I_4) \rangle & \text{if } n \geq 5 \end{cases}$$

(see Example 2.1 for an explicit description of the ideals in this chain). Computations with Macaulay2 [14] yield the following table:

n	4	5	6	7	8	9	10
$\text{pd}(R_n/I_n)$	3	6	8	10	12	14	16

This table suggests that $\text{pd}(R_n/I_n)$ could be a linear function with slope 2 when $n \geq 5$.

The previous example and many other computational experiments lead us to the following expectation:

Conjecture 1.3. *Let $(I_n)_{n \geq 1}$ be a Sym-invariant or, more generally, an Inc^i -invariant chain of ideals. Then $\text{pd}(R_n/I_n)$ is eventually a linear function, that is,*

$$\text{pd}(R_n/I_n) = an + b \quad \text{for some integer constants } a, b \text{ whenever } n \gg 0.$$

It should be noted that this conjecture as well as [21, Conjecture 1.1] is seemingly of parallel nature to the well-known asymptotic linearity of the Castelnuovo-Mumford regularity of the powers of a graded ideal shown independently by Cutkosky, Herzog, and Trung [9, Theorem 1.1(ii)] and Kodiyalam [19, Theorem 5].

Since $\text{codim } I_n$ is eventually a linear function, Conjecture 1.3 is clearly true if I_n is perfect for $n \gg 0$ (see Proposition 4.1). Note that, for example, a graded Cohen-Macaulay ideal is perfect.

It is not hard to give linear upper and lower bounds for $\text{pd}(R_n/I_n)$ (see Proposition 4.3) because

$$cn \geq \text{pd}(R_n/I_n) \geq \text{codim } I_n.$$

Our next main results establish improved lower linear bounds for $\text{pd}(R_n/I_n)$ in the case of a chain of monomial ideals (see Theorems 4.6 and 4.10). These also yield necessary conditions for the Cohen-Macaulayness of R_n/I_n when $n \gg 0$ (see Corollaries 4.7 and 4.12). Note that by the Auslander-Buchsbaum formula all statements on projective dimensions of graded ideals can equivalently be stated as results on depths.

The paper is divided into four sections. Section 2 contains some basic notions and facts on invariant chains of ideals. The asymptotic behavior of codimensions and projective dimensions of ideals in such chains are discussed in Sections 3 and 4, respectively.

Acknowledgement. We are grateful to the anonymous referees for insightful comments and suggestions that helped a lot to improve the clarity of the paper. The first author wishes to thank Lorenzo Venturello for his help with computations using Macaulay2.

2. PRELIMINARIES

We keep the notation and definitions of the introduction. In particular, c is a fixed positive integer, and for each $n \geq 1$, R_n denotes the polynomial ring in $c \times n$ variables over a field K . Let

$$R = \bigcup_{n \geq 1} R_n = K[x_{k,j} \mid 1 \leq k \leq c, j \geq 1]$$

be the polynomial ring in “ $c \times \mathbb{N}$ ” variables. The action of $\text{Sym}(n)$ on R_n given by

$$\sigma \cdot x_{k,j} = x_{k,\sigma(j)} \quad \text{for all } \sigma \in \text{Sym}(n), 1 \leq k \leq c, 1 \leq j \leq n,$$

clearly induces an action of

$$\text{Sym}(\infty) = \bigcup_{n \geq 1} \text{Sym}(n)$$

on R . Recall that a chain of ideals $(I_n)_{n \geq 1}$ with $I_n \subseteq R_n$ is Sym -invariant (or $\text{Sym}(\infty)$ -invariant) if

$$\text{Sym}(n)(I_m) = \{\sigma(f) \mid f \in I_m, \sigma \in \text{Sym}(n)\} \subseteq I_n \quad \text{for all } m \leq n.$$

Often, it is inconvenient to work with Sym -invariant chains. The main reason is that the group $\text{Sym}(\infty)$ is not compatible with monomial orders: For any monomial order \leq on R and any monomials $u, v \in R$ with $u < v$, there exists some $\sigma \in \text{Sym}(\infty)$ such that $\sigma(u) > \sigma(v)$ (see [2, Remark 2.1]). Thus, the initial chain $(\text{in}_{\leq}(I_n))_{n \geq 1}$ of a Sym -invariant chain $(I_n)_{n \geq 1}$ is typically not Sym -invariant (see Example 2.2). Here and in the sequel, whenever \leq is a monomial order on R , we will use the same notation to denote its restrictions to the subrings R_n of R .

To overcome this difficulty, one considers the following monoid of increasing functions on \mathbb{N} :

$$\text{Inc} = \{\pi: \mathbb{N} \rightarrow \mathbb{N} \mid \pi(j) < \pi(j+1) \text{ for all } j \geq 1\},$$

and more generally, submonoids of Inc that fix initial segments of \mathbb{N} :

$$\text{Inc}^i = \{\pi \in \text{Inc} \mid \pi(j) = j \text{ for all } j \leq i\},$$

where $i \geq 0$ is an integer. Observe that one has a descending chain of monoids

$$\text{Inc} = \text{Inc}^0 \supset \text{Inc}^1 \supset \text{Inc}^2 \supset \cdots.$$

The action of Inc^i on R is defined analogously to that of $\text{Sym}(\infty)$. We say that a chain $(I_n)_{n \geq 1}$ with I_n an ideal in R_n is Inc^i -invariant if

$$\text{Inc}_{m,n}^i(I_m) \subseteq I_n \quad \text{for all } m \leq n,$$

where

$$\text{Inc}_{m,n}^i = \{\pi \in \text{Inc}^i \mid \pi(m) \leq n\}.$$

It is evident that every Inc^i -invariant chain is also Inc^{i+1} -invariant. Moreover, for any $f \in R_m$ and $\pi \in \text{Inc}_{m,n}^i$ ($m \leq n$), it is easy to find a permutation $\sigma \in \text{Sym}(n)$ such that $\pi f = \sigma f$ (see, e.g., [22, Lemma 7.6]). Hence, $\text{Inc}_{m,n}^i \cdot f \subseteq \text{Sym}(n) \cdot f$. It follows that every Sym -invariant chain is also an Inc^i -invariant chain.

A fundamental result of Hillar and Sullivant [17, Theorem 3.1] (see also [22, Corollary 3.6]) implies that every Inc^i -invariant chain $\mathcal{J} = (I_n)_{n \geq 1}$ stabilizes, meaning that there exists an integer $r \geq 1$ such that, as ideals in R_n , one has

$$I_n = \langle \text{Inc}_{r,n}^i(I_r) \rangle_{R_n} \quad \text{for all } n \geq r,$$

or equivalently,

$$I_n = \langle \text{Inc}_{m,n}^i(I_m) \rangle_{R_n} \quad \text{for all } n \geq m \geq r$$

(see [22, Lemma 5.2, Corollary 5.4]). The least integer r with this property is called the *i-stability index* of \mathcal{J} , denoted by

$$\text{ind}^i(\mathcal{J}).$$

Example 2.1. Let $\mathcal{J} = (I_n)_{n \geq 1}$ be the Inc^1 -invariant chain considered in Example 1.2. Evidently, $\text{ind}^1(\mathcal{J}) = 4$. Since $x_{2,1}$ is fixed under the action of Inc^1 , some non-zero ideals of \mathcal{J} are

$$\begin{aligned} I_4 &= \langle x_{1,2}^3, x_{1,4}^2 x_{2,1}, x_{2,2} x_{3,3} \rangle, \\ I_5 &= I_4 + \langle x_{1,3}^3, x_{1,5}^2 x_{2,1}, x_{2,2} x_{3,4}, x_{2,3} x_{3,4} \rangle, \\ I_6 &= I_5 + \langle x_{1,4}^3, x_{1,6}^2 x_{2,1}, x_{2,2} x_{3,5}, x_{2,3} x_{3,5}, x_{2,4} x_{3,5} \rangle, \\ I_7 &= I_6 + \langle x_{1,5}^3, x_{1,7}^2 x_{2,1}, x_{2,2} x_{3,6}, x_{2,3} x_{3,6}, x_{2,4} x_{3,6}, x_{2,5} x_{3,6} \rangle. \end{aligned}$$

By induction one can show that for all $n \geq 5$:

$$I_n = I_{n-1} + \langle x_{1,n-2}^3, x_{1,n}^2 x_{2,1}, x_{2,2} x_{3,n-1}, x_{2,3} x_{3,n-1}, \dots, x_{2,n-2} x_{3,n-1} \rangle.$$

When working with invariant chains of ideals, a key advantage of the monoids Inc^i over the group $\text{Sym}(\infty)$ is that the monoids Inc^i behave well with certain monomial orders on R , and the initial chain of any Inc^i -invariant chain with respect to such an order is again Inc^i -invariant (see Lemma 2.3 below). We say that a monomial order \leq *respects* Inc^i if $\pi(u) \leq \pi(v)$ whenever $\pi \in \text{Inc}^i$ and u, v are monomials of R with $u \leq v$. This condition implies that

$$\text{in}_{\leq}(\pi(f)) = \pi(\text{in}_{\leq}(f)) \quad \text{for all } f \in R \text{ and } \pi \in \text{Inc}^i.$$

Examples of monomial orders respecting Inc^i include the lexicographic order and the reverse-lexicographic order on R induced by the following ordering of the variables:

$$x_{k,j} \leq x_{k',j'} \quad \text{if either } k < k' \text{ or } k = k' \text{ and } j < j'.$$

Example 2.2. Let $c = 1$ and consider the ideals

$$\begin{aligned} I_3 &= \langle x_1^2 + x_2x_3, x_2^2 + x_1x_3, x_3^2 + x_1x_2 \rangle, \\ I_4 &= \text{Sym}(4)(I_3) \\ &= I_3 + \langle x_1^2 + x_2x_4, x_1^2 + x_3x_4, x_2^2 + x_1x_4, x_2^2 + x_3x_4, \\ &\quad x_3^2 + x_1x_4, x_3^2 + x_2x_4, x_4^2 + x_1x_2, x_4^2 + x_1x_3, x_4^2 + x_2x_3 \rangle. \end{aligned}$$

Using the reverse-lexicographic order with $x_1 < x_2 < x_3 < x_4$ one obtains by computations with Macaulay2 that

$$\begin{aligned} \text{in}(I_3) &= \langle x_2^2, x_3x_2, x_3^2, x_2x_1^2, x_3x_1^2, x_1^4 \rangle, \\ \text{in}(I_4) &= \langle x_2x_1, x_3x_1, x_4x_1, x_2^2, x_3x_2, x_4x_2, x_3^2, x_4x_3, x_4^2, x_1^3 \rangle. \end{aligned}$$

Since $x_2^2 \in \text{in}(I_3)$, we get $x_1^2 \in \text{Sym}(4)(\text{in}(I_3))$. Thus,

$$\text{Sym}(4)(\text{in}(I_3)) \not\subseteq \text{in}(I_4).$$

Note, however, that

$$\text{Inc}_{3,4}(\text{in}(I_3)) = \text{in}(I_3) + \langle x_4x_2, x_4x_3, x_4^2, x_3x_1^2, x_3x_2^2, x_4x_1^2, x_4x_2^2 \rangle \subseteq \text{in}(I_4).$$

The phenomenon in the preceding example holds true more generally:

Lemma 2.3 ([22, Lemma 7.1]). *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of ideals. Then for any monomial order \leq respecting Inc^i , the chain $\text{in}_{\leq}(\mathcal{J}) = (\text{in}_{\leq}(I_n))_{n \geq 1}$ is also Inc^i -invariant and*

$$\text{ind}^i(\mathcal{J}) \leq \text{ind}^i(\text{in}_{\leq}(\mathcal{J})).$$

We conclude this section with two auxiliary results that will be used frequently. The first one slightly generalizes one part of [22, Lemma 6.4].

Lemma 2.4. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals and $v \in R_i$ a monomial. Then the chain $\mathcal{J} : v = (I_n : v)_{n \geq 1}$ is also Inc^i -invariant and*

$$\text{ind}^i(\mathcal{J} : v) \leq \text{ind}^i(\mathcal{J}).$$

Proof. Write $v = x_{k_1, j_1}^{e_1} \cdots x_{k_m, j_m}^{e_m}$ with $j_1, \dots, j_m \leq i$. By induction on m , we may assume $v = x_{k_1, j_1}^{e_1}$. But in this case the result follows by using the same argument as in the proof of [22, Lemma 6.4]. \square

For the next result we need further notation. The i -shift $\sigma_i \in \text{Inc}^i$ is given by

$$\sigma_i(j) = \begin{cases} j & \text{if } 1 \leq j \leq i, \\ j+1 & \text{if } j \geq i+1. \end{cases}$$

For a graded ideal J in R_n we write $\delta(J)$ for the largest degree of a minimal homogeneous generator of J . Then the q -invariant of J is defined as

$$q(J) = \sum_{j=0}^{\delta(J)} \dim_K(R_n/J)_j.$$

Lemma 2.5. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. For each $\mathbf{e} = (e_1, \dots, e_c) \in \mathbb{Z}_{\geq 0}^c$, consider a chain of monomial ideals $\mathcal{J}_{\mathbf{e}} = (I_{\mathbf{e},n})_{n \geq 1}$ given by*

$$I_{\mathbf{e},n} = \langle (I_n : x_{1,i+1}^{e_1} \cdots x_{c,i+1}^{e_c}), x_{1,i+1}, \dots, x_{c,i+1} \rangle \quad \text{for all } n \geq 1.$$

Then the following statements hold:

- (i) $\mathcal{J}_{\mathbf{e}}$ is an Inc^{i+1} -invariant chain with $\text{ind}^{i+1}(\mathcal{J}_{\mathbf{e}}) \leq \text{ind}^i(\mathcal{J}) + 1$.
- (ii) Fix $\mathbf{e} \in \mathbb{Z}_{\geq 0}^c$. Then for every $r \geq \text{ind}^i(\mathcal{J})$ one has

$$q(I_{\mathbf{e},r+1}) \leq q(I_r),$$

and equality holds if and only if

$$I_{\mathbf{e},n+1} = \langle \sigma_i(I_n), x_{1,i+1}, \dots, x_{c,i+1} \rangle \quad \text{and} \quad R_{n+1}/I_{\mathbf{e},n+1} \cong R_n/I_n \quad \text{for all } n \geq r.$$

Proof. See [21, Lemma 4.3] (and also the proofs of [22, Theorem 6.2, Lemma 6.10, Lemma 6.11]). \square

3. CODIMENSION UP TO SYMMETRY

Fix a nonnegative integer i . From [22, Theorem 7.10] it follows that the codimensions (i.e. heights) of graded ideals in an Inc^i -invariant chain grow eventually linearly. In this section we extend this result to linearity of the codimension of not necessarily graded ideals in an Inc^i -invariant chain. Moreover, the arguments produce an explicit description for the leading coefficient of the linear function.

We first introduce a function that is used to define that leading coefficient. Write $[c] = \{1, \dots, c\}$. For a monomial $1 \neq u \in R_n$, let $\min(u)$ (respectively, $\max(u)$) denote the smallest (respectively, largest) index j such that $x_{k,j}$ divides u for some $k \in [c]$. When J is a proper monomial ideal in R_n with minimal set of monomial generators $G(J)$, we set

$$\begin{aligned} G_i^+(J) &= \{u \in G(J) \mid \min(u) > i\}, \\ G_i(J) &= \{u \in G(J) \mid \min(u) \leq i < \max(u)\}, \\ G_i^-(J) &= \{u \in G(J) \mid \max(u) \leq i\}. \end{aligned}$$

Definition 3.1. Let C be a subset of $[c]$, $u \in R_n$ a monomial, and $J \subsetneq R_n$ a monomial ideal. We say that

- (i) C covers u if there exists $k \in C$ such that $x_{k,j}$ divides u for some $j \geq 1$,
- (ii) C is an i -cover of J if C covers every element of $G_i^+(J)$.

Let

$$\gamma_i(J) = \min\{\#C \mid C \text{ is an } i\text{-cover of } J\}.$$

Note that $0 \leq \gamma_i(J) \leq c$. If $J = R_n$, we adopt the convention that

$$\gamma_i(R_n) = \infty.$$

Example 3.2. Assume $c \geq 3$ and consider the ideal

$$J = \langle x_{2,1}^4, x_{1,1}^3 x_{2,3}^2 x_{1,4}, x_{3,2} x_{1,3}^2 x_{2,4}, x_{2,3}^3 x_{1,4}^2, x_{2,4}^2 x_{3,5}^4 \rangle \subset R_6.$$

Then $G_2^-(J) = \{x_{2,1}^4\}$, $G_2(J) = \{x_{1,1}^3 x_{2,3}^2 x_{1,4}, x_{3,2} x_{1,3}^2 x_{2,4}\}$, $G_2^+(J) = \{x_{2,3}^3 x_{1,4}^2, x_{2,4}^2 x_{3,5}^4\}$. One sees that J has two minimal 2-covers: $C_1 = \{2\}$ and $C_2 = \{1, 3\}$. Thus,

$$\gamma_2(J) = 1.$$

Let us now discuss some basic properties of the function γ_i . For a monomial ideal J we first show that $\gamma_i(J)$ can be computed from a primary decomposition of the ideal $\langle G_i^+(J) \rangle$ (or more efficiently, from the minimal primes of $\langle G_i^+(J) \rangle$). Consider the map φ that assigns the variable $x_{k,j}$ to k for every $j \geq 1$. Then φ clearly induces a map, still denoted by φ , from the set $\text{Min}(\langle G_i^+(J) \rangle)$ of minimal primes of $\langle G_i^+(J) \rangle$ to the set of i -covers of J . Let $\mathcal{C}_i(J)$ be the image of this map, i.e.

$$\mathcal{C}_i(J) = \{\varphi(P) \mid P \in \text{Min}(\langle G_i^+(J) \rangle)\}.$$

Proposition 3.3. Let $J \subsetneq R_n$ be a monomial ideal. If C is a minimal i -cover of J , then $C \in \mathcal{C}_i(J)$. Furthermore,

$$\gamma_i(J) = \min\{\#C \mid C \in \mathcal{C}_i(J)\} = \min\{\#\varphi(P) \mid P \in \text{Min}(\langle G_i^+(J) \rangle)\}.$$

Proof. The first assertion implies the claimed formula for $\gamma_i(J)$ because it says that $\mathcal{C}_i(J)$, which is a subset of the set of i -covers of J , contains all minimal i -covers. So it suffices to prove this assertion. Suppose C is a minimal i -cover of J . Then for each $u \in G_i^+(J)$ there exist $k(u) \in C$ and $j(u) \geq 1$ such that $x_{k(u),j(u)}$ divides u . Set

$$Q = \langle x_{k(u),j(u)} \mid u \in G_i^+(J) \rangle.$$

Then Q is evidently a prime ideal containing $\langle G_i^+(J) \rangle$. It follows that $Q \supseteq P$ for some $P \in \text{Min}(\langle G_i^+(J) \rangle)$. One has

$$C \supseteq \varphi(Q) \supseteq \varphi(P).$$

Due to the minimality of C , this yields $C = \varphi(P) \in \mathcal{C}_i(J)$, because C and $\varphi(P)$ are both i -covers of J . \square

Example 3.4. Consider again the ideal J in Example 3.2. The set of minimal primes of the ideal $\langle G_2^+(J) \rangle = \langle x_{2,3}^3 x_{1,4}^2, x_{2,4}^2 x_{3,5}^4 \rangle$ is

$$\text{Min}(\langle G_2^+(J) \rangle) = \{\langle x_{2,3}, x_{3,5} \rangle, \langle x_{2,3}, x_{2,4} \rangle, \langle x_{1,4}, x_{2,4} \rangle, \langle x_{1,4}, x_{3,5} \rangle\}.$$

Thus,

$$\mathcal{C}_2(J) = \{\{2, 3\}, \{2\}, \{1, 2\}, \{1, 3\}\},$$

and again we find that

$$\gamma_2(J) = \min\{\#C \mid C \in \mathcal{C}_2(J)\} = 1.$$

Some further properties of the function γ_i are given in the following lemmas.

Lemma 3.5. *Let $C \subseteq [c]$, let $u, v \in R_n$ be monomials with $u|v$, and let $J \subseteq J' \subsetneq R_n$ be monomial ideals. Then the following statements hold:*

- (i) *If C covers u , then C also covers v .*
- (ii) *If C is an i -cover of J' , then C is also an i -cover of J .*
- (iii) $\gamma_i(J) \leq \gamma_i(J')$.
- (iv) $\gamma_i(J) = \gamma_i(\sqrt{J})$.

Proof. (i) follows immediately from Definition 3.1(i). Since $J \subseteq J'$, every element of $G_i^+(J)$ is divisible by some element of $G_i^+(J')$. So (ii) is a consequence of (i). From (ii) we get (iii). For (iv) it suffices to show that any i -cover C of J is also an i -cover of \sqrt{J} . Let $v \in G_i^+(\sqrt{J})$. Then v^k is divisible by an element $u \in G_i^+(J)$ for some $k \geq 1$. Since C covers u , it also covers v^k . Hence, C covers v , as desired. Note that one can also prove (iv) by using Proposition 3.3 and the fact that $\langle G_i^+(\sqrt{J}) \rangle = \sqrt{\langle G_i^+(J) \rangle}$. \square

Lemma 3.6. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. Then*

$$\gamma_i(I_n) = \gamma_i(I_{n+1}) \quad \text{for all } n \geq \text{ind}^i(\mathcal{J}).$$

Proof. It suffices to consider proper ideals. We have $\gamma_i(I_n) \leq \gamma_i(I_{n+1})$ for all $n \geq 1$ by Lemma 3.5(iii). For $n \geq \text{ind}^i(\mathcal{J})$ one has $I_{n+1} = \langle \text{Inc}_{n,n+1}^i(I_n) \rangle$, which implies

$$G_i^+(I_{n+1}) \subseteq \text{Inc}_{n,n+1}^i(G_i^+(I_n)).$$

It follows that any i -cover of I_n is also an i -cover of I_{n+1} , because the action of Inc^i keeps the first index of the variables unchanged. Therefore, $\gamma_i(I_n) \geq \gamma_i(I_{n+1})$, and hence $\gamma_i(I_n) = \gamma_i(I_{n+1})$. \square

We set

$$\gamma_i(\mathcal{J}) = \gamma_i(I_n) \quad \text{for some } n \geq \text{ind}^i(\mathcal{J}).$$

This is well-defined by Lemma 3.6, and moreover, according to Proposition 3.3, $\gamma_i(\mathcal{J})$ can be determined by the minimal primes of the ideal $\langle G_i^+(I_n) \rangle$ for any $n \geq \text{ind}^i(\mathcal{J})$.

For convenience in stating and proving the next result we will make use of the following convention:

Convention 3.7. The codimension of the unit ideal in the ring R_n is set to be ∞ for every $n \geq 1$.

The main result of this section is:

Theorem 3.8. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. Then there exists an integer $D(\mathcal{J})$ such that*

$$\text{codim } I_n = \gamma_i(\mathcal{J})n + D(\mathcal{J}) \quad \text{for } n \gg 0.$$

The argument requires some further preparations. The following observation says that it suffices to prove the theorem for chains of squarefree monomial ideals.

Lemma 3.9. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. Then the chain $\sqrt{\mathcal{J}} = (\sqrt{I_n})_{n \geq 1}$ is also Inc^i -invariant with*

$$\gamma_i(\sqrt{\mathcal{J}}) = \gamma_i(\mathcal{J}).$$

Proof. Again it suffices to consider proper ideals. Let $n \geq m \geq 1$, $\pi \in \text{Inc}_{m,n}^i$, and consider any monomial $u \in \sqrt{I_m}$. Let $k \geq 1$ be such that $u^k \in I_m$. Then

$$\pi(u)^k = \pi(u^k) \in \pi(I_m) \subseteq I_n.$$

Thus, $\pi(u) \in \sqrt{I_n}$, and so the chain $\sqrt{\mathcal{J}}$ is Inc^i -invariant. The equality

$$\gamma_i(\sqrt{\mathcal{J}}) = \gamma_i(\mathcal{J})$$

follows from Lemma 3.5(iv). \square

Lemma 3.10. *Let $J \subseteq R_n$ be a squarefree monomial ideal and x a variable of R_n . Then*

$$\text{codim} J = \min\{\text{codim}\langle(J : x), x\rangle - 1, \text{codim}\langle J, x\rangle\}.$$

Proof. If $J = R_n$ or $J : x = R_n$, then the formula is true according to Convention 3.7. If J and $J : x$ are both proper ideals of R_n , then it is apparent that

$$\text{codim} J = \min\{\text{codim}\langle J : x\rangle, \text{codim}\langle J, x\rangle\}.$$

Since J is squarefree, x is a non-zero-divisor on $R_n/\langle J : x\rangle$. This gives

$$\text{codim}\langle J : x\rangle = \text{codim}\langle(J : x), x\rangle - 1,$$

which yields the desired conclusion. \square

The next lemma plays a crucial role in the proof of Theorem 3.8.

Lemma 3.11. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. Fix an integer $r \geq \text{ind}^i(\mathcal{J})$. For each $\mathbf{e} = (e_1, \dots, e_c) \in \mathbb{Z}_{\geq 0}^c$, define the chain $\mathcal{J}_{\mathbf{e}} = (I_{\mathbf{e},n})_{n \geq 1}$ as in Lemma 2.5. Then the following statements hold:*

(i) *If $q(I_{\mathbf{e},r+1}) = q(I_r)$, then*

$$\text{codim} I_{\mathbf{e},n+1} = \text{codim} I_n + c \quad \text{for all } n \geq r.$$

(ii) *If I_n is a squarefree ideal, then*

$$\text{codim} I_n = \min\{\text{codim} I_{\mathbf{e},n} - |\mathbf{e}| \mid \mathbf{e} \in \{0, 1\}^c\},$$

where $|\mathbf{e}| = e_1 + \dots + e_c$.

(iii) *For all $\mathbf{e} \in \mathbb{Z}_{\geq 0}^c$ one has $\gamma_i(\mathcal{J}) \leq \gamma_{i+1}(\mathcal{J}_{\mathbf{e}})$, with equality if $q(I_{\mathbf{e},r+1}) = q(I_r)$.*

(iv) *Assume $I_r \neq R_r$. If $\mathbf{e} \in \{0, 1\}^c$ and $q(I_{\mathbf{e},r+1}) = q(I_r)$, then $\gamma_i(\mathcal{J}) \leq c - |\mathbf{e}|$.*

(v) *Assume $I_r \neq R_r$. Set*

$$E_1 = \{\mathbf{e} \in \{0, 1\}^c \mid q(I_{\mathbf{e},r+1}) < q(I_r)\} \text{ and } E_2 = \{\mathbf{e} \in \{0, 1\}^c \mid q(I_{\mathbf{e},r+1}) = q(I_r)\}.$$

Then

$$(1) \quad \gamma_i(\mathcal{J}) = \min\{\min\{\gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \mid \mathbf{e} \in E_1\}, \min\{c - |\mathbf{e}| \mid \mathbf{e} \in E_2\}\}.$$

Proof. (i) From Lemma 2.5(ii) one gets the isomorphisms

$$R_{n+1}/I_{\mathbf{e},n+1} \cong R_n/I_n \quad \text{for all } n \geq r,$$

which yield the assertion.

(ii) Using Lemma 3.10, the assertion follows by induction on c .

(iii) Let σ_i be the i -shift defined preceding Lemma 2.5. Since $\sigma_i \in \text{Inc}_{n,n+1}^i$, one has

$$I_{\mathbf{e},n+1} \supseteq \langle I_{n+1}, x_{1,i+1}, \dots, x_{c,i+1} \rangle \supseteq \langle \sigma_i(I_n), x_{1,i+1}, \dots, x_{c,i+1} \rangle \quad \text{for all } n \geq 1.$$

By Lemma 3.5(iii), this gives

$$\gamma_{i+1}(I_{\mathbf{e},n+1}) \geq \gamma_{i+1}(\langle \sigma_i(I_n), x_{1,i+1}, \dots, x_{c,i+1} \rangle) = \gamma_{i+1}(\langle \sigma_i(I_n) \rangle) = \gamma_i(I_n).$$

The last equality follows from the definition of σ_i . The only inequality in the above equation becomes an equality if $n \geq r$ and $q(I_{\mathbf{e},r+1}) = q(I_r)$, by Lemma 2.5(ii).

(iv) Set $C = \{k \in [c] \mid e_k = 0\}$. By (iii), it suffices to show that C is an $(i+1)$ -cover of $I_{\mathbf{e},n+1}$ for $n \geq r$. Assume the contrary. Then there exists a monomial $u \in G_{i+1}^+(I_{\mathbf{e},n+1})$ of lowest degree which is not divisible by $x_{k,j}$ for any $k \in C$ and $j \geq 1$. If there is more than one such monomial we choose u with $\min(u)$ as small as possible. Note that $\min(u) \geq i+2$ since $u \in G_{i+1}^+(I_{\mathbf{e},n+1})$. By Lemma 2.5(ii),

$$I_{\mathbf{e},n+1} = \langle \sigma_i(I_n), x_{1,i+1}, \dots, x_{c,i+1} \rangle.$$

It follows that $u \in \sigma_i(I_n)$. So $u = \sigma_i(v)$ for some $v \in I_n$. By definition of σ_i one has

$$\min(v) = \min(u) - 1 \geq i+1.$$

By the choice of u , the monomial v is a minimal generator of $I_{\mathbf{e},n+1}$. If $\min(v) \geq i+2$, then $v \in G_{i+1}^+(I_{\mathbf{e},n+1})$. Since $\min(v) < \min(u)$, this is a contradiction to $\min(u)$ being least possible among all the monomials in $G_{i+1}^+(I_{\mathbf{e},n+1})$ of lowest degree which are not divisible by $x_{k,j}$ for any $k \in C$ and $j \geq 1$. Hence, $\min(v) = i+1$, and we may write $v = x_{l_1,i+1} \cdots x_{l_s,i+1} v'$, where $v' \in R_n$ with $\min(v') > i+1$. Since $u = \sigma_i(v)$ is not divisible by any $x_{k,j}$ with $k \in C$, we must have $l_1, \dots, l_s \notin C$. Thus, $e_{l_1} = \cdots = e_{l_s} = 1$, and so

$$v' \in I_n : x_{l_1,i+1} \cdots x_{l_s,i+1} \subseteq I_{\mathbf{e},n}.$$

Since $\mathcal{J}_{\mathbf{e}}$ is Inc^{i+1} -invariant (see Lemma 2.5) and $\sigma_{i+1} \in \text{Inc}_{n,n+1}^{i+1}$, we obtain $\sigma_{i+1}(v') \in I_{\mathbf{e},n+1}$. As $\min(v') > i+1$, one has $\sigma_i(v') = \sigma_{i+1}(v')$. Thus, $\sigma_i(v') \in I_{\mathbf{e},n+1}$. But this contradicts our assumption that $u = \sigma_i(v) = x_{l_1,i+2} \cdots x_{l_s,i+2} \sigma_i(v')$ is a minimal generator of $I_{\mathbf{e},n+1}$.

(v) Let γ denote the right-hand side of Equation (1). From (iii) and (iv) it follows that $\gamma(\mathcal{J}) \leq \gamma$. For the reverse inequality it suffices to find a tuple $\mathbf{e} \in \{0,1\}^c$ with

$$\gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \leq \gamma(\mathcal{J}) = c - |\mathbf{e}|.$$

Let $n \geq r+1$ and C an i -cover of I_n with $\gamma(I_n) = |C|$. Consider $\mathbf{e} = (e_1, \dots, e_c)$ with

$$e_k = \begin{cases} 0 & \text{if } k \in C, \\ 1 & \text{if } k \notin C. \end{cases}$$

Then it is clear that $\gamma(I_n) = |C| = c - |\mathbf{e}|$. To complete the proof we will show that C is an $(i+1)$ -cover of $I_{\mathbf{e},n}$. For any $u \in G_{i+1}^+(I_{\mathbf{e},n})$ one has

$$v = x_{1,i+1}^{e_1} \cdots x_{c,i+1}^{e_c} u = \prod_{k \notin C} x_{k,i+1} u \in I_n.$$

This implies that v has a divisor $v' \in G_i^+(I_n)$. Since C covers v' , it covers v as well. It then follows that C must cover u . Therefore, C is an $(i+1)$ -cover of $I_{\mathbf{e},n}$, as desired. \square

We are now ready to prove Theorem 3.8.

Proof of Theorem 3.8. Using Lemma 3.9, we may assume that \mathcal{J} is a chain of squarefree monomial ideals. Let \mathcal{F} denote the family of all (i, r, \mathcal{J}) , where $i, r \geq 0$ are integers and $\mathcal{J} = (I_n)_{n \geq 1}$ is an Inc^i -invariant chain of squarefree monomial ideals with $\text{ind}^i(\mathcal{J}) \leq r$.

Following the idea of the proofs of [22, Theorem 6.2] and [21, Theorem 6.2], we argue by induction on $q = q(I_r)$ that for any $(i, r, \mathcal{J}) \in \mathcal{F}$ one has that

$$\text{codim} I_{n+1} = \text{codim} I_n + \gamma_i(\mathcal{J}) \quad \text{whenever } n \gg 0.$$

If $q = 0$, then $I_r = R_r$, and so $I_n = R_n$ for every $n \geq r$. By Convention 3.7, this means that $\text{codim} I_n = \infty$ for $n \geq r$, and the desired conclusion holds according to our convention in Definition 3.1.

Now assume $q \geq 1$. For each $\mathbf{e} \in \{0, 1\}^c$, we consider the chain $\mathcal{J}_{\mathbf{e}} = (I_{\mathbf{e}, n})_{n \geq 1}$ as in Lemma 2.5. By this lemma, $\mathcal{J}_{\mathbf{e}}$ is an Inc^{i+1} -invariant chain with $\text{ind}^{i+1}(\mathcal{J}_{\mathbf{e}}) \leq r + 1$. Write $\{0, 1\}^c = E_1 \cup E_2$ with

$$E_1 = \{\mathbf{e} \in \{0, 1\}^c \mid q(I_{\mathbf{e}, r+1}) < q\} \quad \text{and} \quad E_2 = \{\mathbf{e} \in \{0, 1\}^c \mid q(I_{\mathbf{e}, r+1}) = q\}.$$

If $\mathbf{e} \in E_2$, then Lemma 3.11(i) gives

$$\text{codim} I_{\mathbf{e}, n} = \text{codim} I_{n-1} + c \quad \text{for all } n \geq r + 1.$$

Hence, for all $n \geq r + 1$ it follows from Lemma 3.11(ii) that

$$\begin{aligned} \text{codim} I_n &= \min \{ \min \{ \text{codim} I_{\mathbf{e}, n} - |\mathbf{e}| \mid \mathbf{e} \in E_1 \}, \min \{ \text{codim} I_{\mathbf{e}, n} - |\mathbf{e}| \mid \mathbf{e} \in E_2 \} \} \\ (2) \quad &= \min \{ \min \{ \text{codim} I_{\mathbf{e}, n} - |\mathbf{e}| \mid \mathbf{e} \in E_1 \}, \text{codim} I_{n-1} + \min \{ c - |\mathbf{e}| \mid \mathbf{e} \in E_2 \} \}. \end{aligned}$$

For $\mathbf{e} \in E_1$, the induction hypothesis applied to $(i + 1, r + 1, \mathcal{J}_{\mathbf{e}}) \in \mathcal{F}$ yields the existence of an integer $N(\mathcal{J}_{\mathbf{e}}) \geq r + 1$ such that

$$(3) \quad \text{codim} I_{\mathbf{e}, n+1} = \text{codim} I_{\mathbf{e}, n} + \gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \quad \text{whenever } n \geq N(\mathcal{J}_{\mathbf{e}}).$$

Set

$$N = \max \{ N(\mathcal{J}_{\mathbf{e}}) \mid \mathbf{e} \in E_1 \}.$$

We will show

$$(4) \quad \text{codim} I_{n+1} = \text{codim} I_n + \gamma_i(\mathcal{J}) \quad \text{whenever } n \geq N.$$

Indeed, by Lemma 3.11(v)

$$(5) \quad \gamma_i(\mathcal{J}) = \min \{ \min \{ \gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \mid \mathbf{e} \in E_1 \}, \min \{ c - |\mathbf{e}| \mid \mathbf{e} \in E_2 \} \}.$$

If $n \geq N$ and $\mathbf{e} \in E_1$, then

$$\begin{aligned} \text{codim} I_{\mathbf{e}, n+1} - |\mathbf{e}| &= \text{codim} I_{\mathbf{e}, n} + \gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) - |\mathbf{e}| \\ &\geq \text{codim} I_n + \gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) && \text{by Equation (2)} \\ &\geq \text{codim} I_n + \gamma_i(\mathcal{J}) && \text{by Equation (5)}. \end{aligned}$$

Combined with Equations (2) and (5), this implies

$$\begin{aligned} (6) \quad \text{codim} I_{n+1} &= \min \{ \min \{ \text{codim} I_{\mathbf{e}, n+1} - |\mathbf{e}| \mid \mathbf{e} \in E_1 \}, \text{codim} I_n + \min \{ c - |\mathbf{e}| \mid \mathbf{e} \in E_2 \} \} \\ &\geq \min \{ \min \{ \text{codim} I_n + \gamma_i(\mathcal{J}) \mid \mathbf{e} \in E_1 \}, \text{codim} I_n + \min \{ c - |\mathbf{e}| \mid \mathbf{e} \in E_2 \} \} \\ &\geq \text{codim} I_n + \gamma_i(\mathcal{J}) \quad \text{if } n \geq N. \end{aligned}$$

Moreover, Equation (2) gives

$$\text{codim } I_{n+1} \leq \text{codim } I_n + \min\{c - |\mathbf{e}| \mid \mathbf{e} \in E_2\} \quad \text{if } n \geq N.$$

This together with Inequality (6) yields Equation (4) if $\gamma_i(\mathcal{J}) = \min\{c - |\mathbf{e}| \mid \mathbf{e} \in E_2\}$.

Thus, it remains to consider the case $\gamma_i(\mathcal{J}) < \min\{c - |\mathbf{e}| \mid \mathbf{e} \in E_2\}$. Suppose Equation (4) is not true. Taking into account Inequality (6), this means that, for any $n_0 \geq N$, there is some $n > n_0$ with

$$\text{codim } I_{n+1} > \text{codim } I_n + \gamma_i(\mathcal{J}).$$

We use this to define an increasing sequence $(n_j)_{j \in \mathbb{N}}$ of integers: set $n_0 = N$ and, for $j \geq 1$, let n_j be the least integer $n > n_{j-1}$ with $\text{codim } I_{n+1} > \text{codim } I_n + \gamma_i(\mathcal{J})$. Thus, we obtain for every $j \geq 1$,

$$(7) \quad \text{codim } I_{n_j+1} \geq \text{codim } I_N + (n_j + 1 - N)\gamma_i(\mathcal{J}) + j.$$

Our assumption $\gamma_i(\mathcal{J}) < \min\{c - |\mathbf{e}| \mid \mathbf{e} \in E_2\}$ allows us to fix some $\mathbf{e}_0 \in E_1$ such that $\gamma_i(\mathcal{J}) = \gamma_{i+1}(\mathcal{J}_{\mathbf{e}_0})$. Let j be an integer with $j > \text{codim } I_{\mathbf{e}_0, N} - \text{codim } I_N - |\mathbf{e}_0|$, i.e.,

$$\text{codim } I_N + (n_j + 1 - N)\gamma_i(\mathcal{J}) + j > \text{codim } I_{\mathbf{e}_0, N} + (n_j + 1 - N)\gamma_i(\mathcal{J}) - |\mathbf{e}_0|.$$

Combining this with Inequality (7) one gets

$$\begin{aligned} \text{codim } I_{n_j+1} &> \text{codim } I_{\mathbf{e}_0, N} + (n_j + 1 - N)\gamma_i(\mathcal{J}) - |\mathbf{e}_0| \\ &= \text{codim } I_{\mathbf{e}_0, N} + (n_j + 1 - N)\gamma_{i+1}(\mathcal{J}_{\mathbf{e}_0}) - |\mathbf{e}_0| \\ &= \text{codim } I_{\mathbf{e}_0, n_j+1} - |\mathbf{e}_0| \quad \text{by Equation (3).} \end{aligned}$$

However, this contradicts Equation (2). The proof is complete. \square

As a consequence, we obtain an explicit and more general version of the first part of [22, Theorem 7.10]. We use the convention that the dimension of a zero module is $-\infty$.

Corollary 3.12. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of ideals, and let \leq be any monomial order respecting Inc^i . Then one has*

$$\dim R_n / I_n = A(\mathcal{J})n + B(\mathcal{J}) \quad \text{for } n \gg 0,$$

where $A(\mathcal{J}) = c - \gamma_i(\text{in}_{\leq}(\mathcal{J}))$ and $B(\mathcal{J}) = -D(\text{in}_{\leq}(\mathcal{J}))$. In particular, the integers $\gamma_i(\mathcal{J}) = \gamma_i(\text{in}_{\leq}(\mathcal{J}))$ and $D(\mathcal{J}) = D(\text{in}_{\leq}(\mathcal{J}))$ are independent of the choice of \leq .

Proof. Since $\text{codim } I_n = \text{codim}(\text{in}_{\leq}(I_n))$ for all $n \geq 1$ (see, e.g., [3, Proposition 3.1(a)]), the result follows from Lemma 2.3 and Theorem 3.8. \square

Remark 3.13. Let $\mathcal{J} = (I_n)_{n \geq 1}$ be a chain of ideals. If this chain is Inc^i -invariant for some $i \geq 0$, then by definition, it is also Inc^j -invariant for all $j \geq i$. So Corollary 3.12 implies that $\gamma_i(\mathcal{J}) = \gamma_j(\mathcal{J})$ for all $j \geq i$. Hence, $\gamma_i(\mathcal{J})$ is independent of the choice of i such that \mathcal{J} is an Inc^i -invariant chain.

4. PROJECTIVE DIMENSION UP TO SYMMETRY

In this section we provide evidence for Conjecture 1.3. First, recall that an ideal $J \subseteq R_n$ is *perfect* if $\text{codim} J = \text{pd}(R_n/J)$. Theorem 3.8 thus implies:

Proposition 4.1. *If $\mathcal{J} = (I_n)_{n \geq 1}$ is an Inc^i -invariant chain of ideals such that I_n is perfect for all $n \gg 0$, then Conjecture 1.3 is true for \mathcal{J} .*

Example 4.2. Chains that satisfy the assumption of Proposition 4.1 include the following interesting ones:

- (i) \mathcal{J} is generated by one monomial orbit; see [15, Corollary 2.2].
- (ii) I_r is an Artinian ideal in R_r for some $r \geq \text{ind}^i(\mathcal{J})$.
- (iii) I_n is generated by the t -minors of the $c \times n$ matrix with entries being the variables of R_n for all $n \geq 1$, where $t \leq c$ is a fixed integer; see, e.g., [4, Theorem 7.3.1].
- (iv) More generally, \mathcal{J} is a chain of graded ideals such that R_n/I_n is Cohen-Macaulay for $n \gg 0$; see, e.g., [4, Corollary 2.2.15].

Using the notation of Corollary 3.12, it is not hard to give linear bounds for $\text{pd}(R_n/I_n)$:

Proposition 4.3. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper ideals. Then one has*

$$cn \geq \text{pd}(R_n/I_n) \geq \gamma_i(\mathcal{J})n + D(\mathcal{J}) \quad \text{for } n \gg 0.$$

Proof. The upper bound is Hilbert's Syzygy theorem. For the lower bound, note that $I_n \neq R_n$ for all $n \geq 1$ by assumption. By using Corollary 3.12 and the estimate

$$\text{pd}(R_n/I_n) \geq \text{codim} I_n \quad \text{for all } n \geq 1$$

(see, e.g., [5, Corollary 16.12]), the desired conclusion follows. \square

In the remaining part of this section we focus on chains of proper monomial ideals. Our main results are lower linear bounds for $\text{pd} I_n = \text{pd}(R_n/I_n) - 1$ that improve the bound given in Proposition 4.3.

Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper monomial ideals. Fix an integer $r \geq \text{ind}^i(\mathcal{J})$. For each $\mathbf{e} \in \mathbb{Z}_{\geq 0}^c$ consider the chain $\mathcal{J}_{\mathbf{e}} = (I_{\mathbf{e},n})_{n \geq 1}$ as in Lemma 2.5. We know that $\mathcal{J}_{\mathbf{e}}$ is an Inc^{i+1} -invariant chain with $\gamma_i(\mathcal{J}) \leq \gamma_{i+1}(\mathcal{J}_{\mathbf{e}})$. Let

$$(8) \quad E(\mathcal{J}) = \{\mathbf{e} \in \mathbb{Z}_{\geq 0}^c \mid I_{\mathbf{e},r+1} \neq R_{r+1}\}.$$

Define

$$\gamma_{i+1}^{\max}(\mathcal{J}) = \max\{\gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \mid \mathbf{e} \in E(\mathcal{J})\}.$$

Similarly, for each $\mathbf{e} \in E(\mathcal{J})$ and $\mathbf{e}' = (e'_1, \dots, e'_c) \in \mathbb{Z}_{\geq 0}^c$ one can build an Inc^{i+2} -invariant chain $\mathcal{J}_{\mathbf{e},\mathbf{e}'} = (I_{\mathbf{e},\mathbf{e}',n})_{n \geq 1}$ with

$$I_{\mathbf{e},\mathbf{e}',n} = \langle (I_{\mathbf{e},n} : x_{1,i+2}^{e'_1} \cdots x_{c,i+2}^{e'_c}), x_{1,i+2}, \dots, x_{c,i+2} \rangle \quad \text{for all } n \geq 1.$$

Then define

$$E(\mathcal{J}_{\mathbf{e}}) = \{\mathbf{e}' \in \mathbb{Z}_{\geq 0}^c \mid I_{\mathbf{e},\mathbf{e}',r+2} \neq R_{r+2}\}$$

and

$$\gamma_{i+2}^{\max}(\mathcal{J}) = \max\{\gamma_{i+2}(\mathcal{J}_{\mathbf{e},\mathbf{e}'}) \mid \mathbf{e} \in E(\mathcal{J}), \mathbf{e}' \in E(\mathcal{J}_{\mathbf{e}})\}.$$

Repeating this construction we obtain a non-decreasing sequence of integers

$$\gamma_i(\mathcal{J}) \leq \gamma_{i+1}^{\max}(\mathcal{J}) \leq \gamma_{i+2}^{\max}(\mathcal{J}) \leq \dots \leq c.$$

Let $\Gamma_i(\mathcal{J})$ denote the limit of this sequence:

$$\Gamma_i(\mathcal{J}) = \max\{\gamma_{i+k}^{\max}(\mathcal{J}) \mid k \geq 1\}.$$

It is obvious that

$$\Gamma_i(\mathcal{J}) \geq \gamma_i(\mathcal{J}).$$

This inequality is strict in general, as illustrated below.

Example 4.4. Let $r \geq i + 2$ and consider the chain $\mathcal{J} = (I_n)_{n \geq 1}$ with

$$I_n = \begin{cases} \langle 0 \rangle & \text{if } n < r, \\ \langle x_{1,i+1}x_{1,i+2}, x_{1,i+1}x_{2,i+2}, \dots, x_{1,i+1}x_{c,i+2} \rangle & \text{if } n = r, \\ \langle \text{Inc}_{r,n}^i(I_r) \rangle & \text{if } n > r. \end{cases}$$

Then $\text{ind}^i(\mathcal{J}) = r$. One has $\gamma_i(\mathcal{J}) = 1$, since $C = \{1\}$ is an i -cover of I_r . Now for $\mathbf{e} = (1, 0, \dots, 0)$ it is easily seen that

$$I_{\mathbf{e}, r+1} = \langle x_{1,i+3}, \dots, x_{c,i+3}, x_{1,i+2}, \dots, x_{c,i+2}, x_{1,i+1}, \dots, x_{c,i+1} \rangle.$$

This gives $\gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) = c$, which implies $\Gamma_i(\mathcal{J}) = c$, because $\gamma_{i+1}(\mathcal{J}_{\mathbf{e}}) \leq \Gamma_i(\mathcal{J}) \leq c$. Hence, if $c > 1$ then

$$c = \Gamma_i(\mathcal{J}) > \gamma_i(\mathcal{J}) = 1,$$

and the difference is as large as possible.

Proposition 3.3 provides an efficient way to compute $\gamma_i(\mathcal{J})$. It would be interesting to know a similar result for $\Gamma_i(\mathcal{J})$.

Question 4.5. How to compute $\Gamma_i(\mathcal{J})$? In particular, with notation of Proposition 3.3, is it true that

$$\Gamma_i(\mathcal{J}) \in \min\{\#C \mid C \in \mathcal{C}_i(I_n)\} \quad \text{for some } n \geq \text{ind}^i(\mathcal{J})?$$

Let us now improve the lower bound in Proposition 4.3 for chains of monomial ideals.

Theorem 4.6. Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper monomial ideals. Then there exists an integer $\tilde{D}(\mathcal{J})$ such that

$$\text{pd } I_n \geq \Gamma_i(\mathcal{J})n + \tilde{D}(\mathcal{J}) \quad \text{for } n \gg 0.$$

This result gives the following necessary condition for eventual Cohen-Macaulayness of Inc^i -invariant chains of monomial ideals:

Corollary 4.7. Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper ideals. If there is a monomial order \leq respecting Inc^i and such that $R_n/\text{in}_{\leq}(I_n)$ is Cohen-Macaulay for $n \gg 0$, then

$$\gamma_i(\mathcal{J}) = \Gamma_i(\text{in}_{\leq}(\mathcal{J})).$$

Proof. Since $R_n/\text{in}_{\leq}(I_n)$ is Cohen-Macaulay if and only if $\text{codim } \text{in}_{\leq}(I_n) = \text{pd } R_n/\text{in}_{\leq}(I_n)$, the result follows by combining Theorems 3.8 and 4.6. \square

To prove Theorem 4.6 we need some auxiliary results.

Lemma 4.8. *Let $J \subseteq R_n$ be a monomial ideal and x a variable of R_n . Then*

$$\max\{\mathrm{pd}\langle J : x \rangle, \mathrm{pd}\langle J, x \rangle - 1\} \leq \mathrm{pd}J \in \{\mathrm{pd}\langle J : x \rangle, \mathrm{pd}\langle J, x \rangle\}.$$

Moreover, if $d \geq 1$ is an integer such that $J : x^d = J : x^{d+1}$, then

$$\max\{\mathrm{pd}\langle (J : x^k), x \rangle \mid 0 \leq k \leq d\} - 1 \leq \mathrm{pd}J \in \{\mathrm{pd}\langle (J : x^d), x \rangle - 1, \mathrm{pd}\langle (J : x^k), x \rangle \mid 0 \leq k < d\}.$$

Proof. The containment in the first assertion follows from [6, Corollary 3.3(i)] and the Auslander-Buchsbaum formula, whereas the lower bound follows from the containment and the exact sequence

$$0 \rightarrow (R_n / \langle J : x \rangle)(-1) \rightarrow R_n / J \rightarrow R_n / \langle J, x \rangle \rightarrow 0.$$

Now applying the first assertion to the ideals $J : x^k$ for $k \geq 0$ we get

$$(9) \quad \max\{\mathrm{pd}\langle J : x^{k+1} \rangle, \mathrm{pd}\langle (J : x^k), x \rangle - 1\} \leq \mathrm{pd}\langle J : x^k \rangle \in \{\mathrm{pd}\langle J : x^{k+1} \rangle, \mathrm{pd}\langle (J : x^k), x \rangle\}.$$

Since $J : x^d = J : x^{d+1}$, x is a non-zero-divisor on $R_n / \langle J : x^d \rangle$, which gives

$$\mathrm{pd}\langle J : x^{d+1} \rangle = \mathrm{pd}\langle J : x^d \rangle = \mathrm{pd}\langle (J : x^d), x \rangle - 1.$$

Combining this with (9) for $k = 0, \dots, d$ yields the second assertion. \square

Lemma 4.9. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper monomial ideals. For each $\mathbf{e} \in \mathbb{Z}_{\geq 0}^c$ consider the chain $\mathcal{J}_{\mathbf{e}} = (I_{\mathbf{e},n})_{n \geq 1}$ as in Lemma 2.5. Let $E(\mathcal{J})$ be defined as in Equation (8). Then*

$$\max\{\mathrm{pd}I_{\mathbf{e},n} \mid \mathbf{e} \in E(\mathcal{J})\} - c \leq \mathrm{pd}I_n \leq \max\{\mathrm{pd}I_{\mathbf{e},n} \mid \mathbf{e} \in E(\mathcal{J})\} \quad \text{for all } n \geq r + 1.$$

Proof. By induction on c it suffices to consider the case $c = 1$. Let $n \geq r + 1$ and choose an integer $d \geq 1$ such that $I_n : x_{1,i+1}^d = I_n : x_{1,i+1}^{d+1}$. Then Lemma 4.8 gives

$$(10) \quad \begin{aligned} \max\{\mathrm{pd}I_{\mathbf{e},n} \mid 0 \leq \mathbf{e} \leq d\} - 1 &= \max\{\mathrm{pd}\langle (I_n : x_{1,i+1}^{\mathbf{e}}), x_{1,i+1} \rangle \mid 0 \leq \mathbf{e} \leq d\} - 1 \\ &\leq \mathrm{pd}I_n \leq \max\{\mathrm{pd}\langle (I_n : x_{1,i+1}^{\mathbf{e}}), x_{1,i+1} \rangle \mid 0 \leq \mathbf{e} \leq d\} = \max\{\mathrm{pd}I_{\mathbf{e},n} \mid 0 \leq \mathbf{e} \leq d\}. \end{aligned}$$

So to complete the proof we need to show that

$$\max\{\mathrm{pd}I_{\mathbf{e},n} \mid 0 \leq \mathbf{e} \leq d\} = \max\{\mathrm{pd}I_{\mathbf{e},n} \mid \mathbf{e} \in E(\mathcal{J})\}.$$

Note that $I_{\mathbf{e},n} = I_{d,n}$ for all $\mathbf{e} \geq d$ since $I_n : x_{1,i+1}^d = I_n : x_{1,i+1}^{d+1}$. On the other hand, $I_{\mathbf{e},n} = R_n$ has projective dimension 0 for all $\mathbf{e} \in \mathbb{Z}_{\geq 0} \setminus E(\mathcal{J})$. Therefore,

$$\max\{\mathrm{pd}I_{\mathbf{e},n} \mid 0 \leq \mathbf{e} \leq d\} = \max\{\mathrm{pd}I_{\mathbf{e},n} \mid \mathbf{e} \in \mathbb{Z}_{\geq 0}\} = \max\{\mathrm{pd}I_{\mathbf{e},n} \mid \mathbf{e} \in E(\mathcal{J})\}. \quad \square$$

Now we prove Theorem 4.6.

Proof of Theorem 4.6. Let $k \geq 1$ be such that $\Gamma_i(\mathcal{J}) = \gamma_{i+k}^{\max}(\mathcal{J})$. Applying Lemma 4.9 iteratively we obtain

$$\mathrm{pd}I_n \geq \max\{\mathrm{pd}I_{\mathbf{e}_1, \dots, \mathbf{e}_k, n} \mid \mathbf{e}_1 \in E(\mathcal{J}), \dots, \mathbf{e}_k \in E(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_{k-1}})\} - kc$$

for all $n \gg 0$. By Theorem 3.8,

$$\mathrm{pd}I_{\mathbf{e}_1, \dots, \mathbf{e}_k, n} \geq \mathrm{codim}I_{\mathbf{e}_1, \dots, \mathbf{e}_k, n} - 1 = \gamma_{i+k}(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_k})n + D(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_k}) - 1 \quad \text{for } n \gg 0.$$

So if we set

$$\tilde{D}(\mathcal{J}) = \min\{D(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_k}) - 1 \mid \mathbf{e}_1 \in E(\mathcal{J}), \dots, \mathbf{e}_k \in E(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_{k-1}})\} - kc,$$

then it follows that

$$\begin{aligned} \text{pd} I_n &\geq \max\{\gamma_{i+k}(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_k}) \mid \mathbf{e}_1 \in E(\mathcal{J}), \dots, \mathbf{e}_k \in E(\mathcal{J}_{\mathbf{e}_1, \dots, \mathbf{e}_{k-1}})\}n + \tilde{D}(\mathcal{J}) \\ &= \gamma_{i+k}^{\max}(\mathcal{J})n + \tilde{D}(\mathcal{J}) = \Gamma_i(\mathcal{J})n + \tilde{D}(\mathcal{J}) \end{aligned}$$

for all $n \gg 0$. \square

Next, we discuss another improvement of the lower bound given in Proposition 4.3. Assume $r \geq \text{ind}^i(\mathcal{J})$ and let $G(I_r)$ be the minimal set of monomial generators of I_r . Evidently, every monomial $u \in G_i(I_r)$ can be uniquely written as $u = u_1 u_2$ with $\max(u_1) \leq i$, $\min(u_2) > i$. Set

$$G_{i,1}(I_r) = \{u_1 \mid u \in G_i(I_r)\} \quad \text{and} \quad G_{i,2}(I_r) = \{u_2 \mid u \in G_i(I_r)\}.$$

Observe that the sets $G_i^-(I_r)$ and $G_{i,1}(I_r)$ are fixed under the action of Inc^i , whereas $G_i^+(I_r)$ and $G_{i,2}(I_r)$ usually change. So intuitively, one would expect that the growth of $\text{pd} I_n$ depends on $G_i^+(I_r)$ and $G_{i,2}(I_r)$. This will be clarified now.

For a subset M of $G_i(I_r)$, set

$$v_M = \prod_{u \in M} u_1 \in R_i$$

and consider the chain

$$\mathcal{J} : v_M = (I_n : v_M)_{n \geq 1}.$$

According to Lemma 2.4, $\mathcal{J} : v_M$ is an Inc^i -invariant chain with $\text{ind}^i(\mathcal{J} : v_M) \leq \text{ind}^i(\mathcal{J})$. Assume that the monomial v_M is not divisible by any element of $G_i^-(I_r)$. Then $I_r : v_M \neq R_r$ and $G_i^+(I_r : v_M)$ consists of minimal elements (under divisibility) of $G_i^+(I_r) \cup \{u_2 \mid u \in M\}$. It follows that

$$\gamma_i(\mathcal{J}) = \gamma_i(\langle G_i^+(I_r) \rangle) \leq \gamma_i(\langle G_i^+(I_r) \cup \{u_2 \mid u \in M\} \rangle) = \gamma_i(\langle G_i^+(I_r : v_M) \rangle) = \gamma_i(\mathcal{J} : v_M).$$

Thus, the following lower bound also improves the one in Proposition 4.3.

Theorem 4.10. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals. Let $r \geq \text{ind}^i(\mathcal{J})$ and denote by \mathcal{M} the set of all subsets M of $G_i(I_r)$ such that v_M is not divisible by any element of $G_i^-(I_r)$. Then there exists an integer $\tilde{D}(\mathcal{J})$ such that*

$$\text{pd} I_n \geq \max\{\gamma_i(\mathcal{J} : v_M) \mid M \in \mathcal{M}\}n + \tilde{D}(\mathcal{J}) \quad \text{for } n \gg 0.$$

Proof. Using the lower bound in Lemma 4.8 repeatedly, one gets $\text{pd} I_n \geq \text{pd}(I_n : v_M)$. \square

Let us briefly compare the bounds in Theorems 4.6 and 4.10. Observe that $\Gamma_i(\mathcal{J})$ only depends on $G_i^+(I_r)$, while the bound in Theorem 4.10 depends on $G_i^+(I_r)$ and $G_{i,2}(I_r)$ (and also on $G_i^-(I_r)$ and $G_{i,1}(I_r)$). So for instance, the bound in Theorem 4.6 is potentially better if $G_i(I_r) = \emptyset$, while the one in Theorem 4.10 is potentially better if $G_i^+(I_r) = \emptyset$.

As an immediate consequence of Theorem 4.10, we obtain the following bound, which depends only on $G_i^+(I_r)$ and $G_{i,2}(I_r)$.

Corollary 4.11. *Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of monomial ideals with $\text{ind}^i(\mathcal{J}) \leq r$. Assume that*

$$v_{G_i(I_r)} = \prod_{u \in G_i(I_r)} u_1$$

is not divisible by any element of $G_i^-(I_r)$. Then there exists a constant $\tilde{D}(\mathcal{J})$ such that

$$\mathrm{pd} I_n \geq \gamma_i(\langle G_i^+(I_r) \cup G_{i,2}(I_r) \rangle) n + \tilde{D}(\mathcal{J}) \quad \text{for } n \gg 0.$$

Note that the assumption of Corollary 4.11 is satisfied if $G_i^-(I_r) = \emptyset$.

Similarly to Corollary 4.7, one gets from Theorem 4.10 another necessary condition for eventual Cohen-Macaulayness of Inc^i -invariant chains. We only state a version for monomial ideals and leave the more general statement to the interested reader.

Corollary 4.12. *With assumption as in Theorem 4.10, if R_n/I_n is Cohen-Macaulay for $n \gg 0$, then*

$$\gamma_i(\mathcal{J}) = \max\{\gamma_i(\mathcal{J} : v_M) \mid M \in \mathcal{M}\}.$$

To conclude this section, we consider the case $c = 1$, i.e., there is only one row of variables. The next result shows that Conjecture 1.3 is “nearly true” in this case.

Proposition 4.13. *Assume $c = 1$. Let $\mathcal{J} = (I_n)_{n \geq 1}$ be an Inc^i -invariant chain of proper monomial ideals. Then either $\mathrm{pd}(R_n/I_n)$ is eventually a constant or there exists a nonnegative integer D such that*

$$n - D \leq \mathrm{pd}(R_n/I_n) \leq n \quad \text{for all } n \gg 0.$$

Proof. Let $r \geq \mathrm{ind}^i(\mathcal{J})$. We distinguish three cases:

Case 1: $G_i^+(I_r) = G_i(I_r) = \emptyset$. In this case, $G(I_r) = G_i^-(I_r)$ is fixed under the action of Inc^i . It follows that

$$I_n = \langle \mathrm{Inc}_{r,n}^i(I_r) \rangle_{R_n} = \langle I_r \rangle_{R_n} \quad \text{for all } n \geq r.$$

Hence, $\mathrm{pd}(R_n/I_n) = \mathrm{pd}(R_r/I_r)$ for all $n \geq r$.

Case 2: $G_i^+(I_r) \neq \emptyset$. Then $\gamma_i(\mathcal{J}) \geq 1$. Since $\gamma_i(\mathcal{J}) \leq c = 1$, we must have $\gamma_i(\mathcal{J}) = 1$. Applying Proposition 4.3 the result follows.

Case 3: $G_i(I_r) \neq \emptyset$. Let $u \in G_i(I_r)$ and write $u = u_1 u_2$ with $\max(u_1) \leq i$, $\min(u_2) > i$. Since u is a minimal generator of I_r , u_1 is not divisible by any element of $G_i^-(I_r)$. Consider the chain $\mathcal{J} : u_1$. One has $\gamma_i(\mathcal{J} : u_1) \geq 1$ since $u_2 \in G_i^+(I_r : u_1)$. So using Proposition 4.3 and Theorem 4.10 concludes the proof. \square

REFERENCES

- [1] M. Aschenbrenner and C.J. Hillar, *Finite generation of symmetric ideals*. Trans. Amer. Math. Soc. **359** (2007), no. 11, 5171–5192.
- [2] A.E. Brouwer and J. Draisma, *Equivariant Gröbner bases and the Gaussian two-factor model*. Math. Comp. **80** (2011), no. 274, 1123–1133.
- [3] W. Bruns and A. Conca, *Gröbner bases, initial ideals and initial algebras*. In: L.L. Avramov et al. (Hrsg.), Homological methods in commutative algebra, IPM Proceedings, Teheran, 2004.
- [4] W. Bruns and J. Herzog, *Cohen-Macaulay rings. Revised edition*. Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1998.
- [5] W. Bruns and U. Vetter, *Determinantal Rings*. Lecture Notes in Mathematics **1327**, Springer-Verlag, 1988.
- [6] G. Caviglia, H.T. Hà, J. Herzog, M. Kummini, N. Terai, and N.V. Trung, *Depth and regularity modulo a principal ideal*. To appear in J. Algebraic Combin. (2018).
- [7] T. Church, J.S. Ellenberg, and B. Farb, *FI-modules and stability for representations of symmetric groups*. Duke Math. J. **164** (2015), no. 9, 1833–1910.

- [8] D.E. Cohen, *On the laws of a metabelian variety*. J. Algebra **5** (1967), 267–273.
- [9] S.D. Cutkosky, J. Herzog, and N.V. Trung, *Asymptotic behaviour of the Castelnuovo-Mumford regularity*. Compositio Math. **118** (1999), no. 3, 243–261.
- [10] J. Draisma, *Finiteness of the k -factor model and chirality varieties*. Adv. Math. **223** (2010), no. 1, 243–256.
- [11] J. Draisma, *Noetherianity up to symmetry*. In: *Combinatorial algebraic geometry*, Lecture Notes in Mathematics **2108**, pp. 33–61, Springer, 2014.
- [12] J. Draisma, R. Eggermont, R. Krone, and A. Leykin, *Noetherianity for infinite-dimensional toric varieties*. Algebra Number Theory **9** (2015), no. 8, 1857–1880.
- [13] J. Draisma and J. Kuttler, *Bounded-rank tensors are defined in bounded degree*. Duke Math. J. **163** (2014), no. 1, 35–63.
- [14] D. Grayson and M. Stillman, *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [15] S. Güntürkün and U. Nagel, *Equivariant Hilbert series of monomial orbits*. Proc. Amer. Math. Soc. **146** (2018), no. 6, 2381–2393.
- [16] C. Hillar and A. Martín del Campo, *Finiteness theorems and algorithms for permutation invariant chains of Laurent lattice ideals*. J. Symbolic Comput. **50** (2013), 314–334.
- [17] C.J. Hillar and S. Sullivant, *Finite Gröbner bases in infinite dimensional polynomial rings and applications*. Adv. Math. **229** (2012), no. 1, 1–25.
- [18] S. Hoşten and S. Sullivant, *A finiteness theorem for Markov bases of hierarchical models*. J. Combin. Theory Ser. A **114** (2007), no. 2, 311–321.
- [19] V. Kodiyalam, *Asymptotic behaviour of Castelnuovo-Mumford regularity*. Proc. Amer. Math. Soc. **128** (2000), no. 2, 407–411.
- [20] R. Krone, A. Leykin, and A. Snowden, *Hilbert series of symmetric ideals in infinite polynomial rings via formal languages*. J. Algebra **485** (2017), 353–362.
- [21] D.V. Le, U. Nagel, H.D. Nguyen, and T. Römer, *Castelnuovo-Mumford regularity up to symmetry*. Preprint, 2018.
- [22] U. Nagel and T. Römer, *Equivariant Hilbert series in non-Noetherian polynomial rings*. J. Algebra **486** (2017), 204–245.
- [23] U. Nagel and T. Römer, *FI- and OI-modules with varying coefficients*. Preprint, 2017, [arXiv:1710.09247](https://arxiv.org/abs/1710.09247).
- [24] A. Putman and S.V. Sam, *Representation stability and finite linear groups*. Duke Math. J. **166** (2017), no. 13, 2521–2598.
- [25] S.V. Sam and A. Snowden, *Gröbner methods for representations of combinatorial categories*. J. Amer. Math. Soc. **30** (2017), no. 1, 159–203.
- [26] S.V. Sam and A. Snowden, *Representations of categories of G -maps*. To appear in J. Reine Angew. Math.
- [27] F. Santos and B. Sturmfels, *Higher Lawrence configurations*. J. Combin. Theory Ser. A **103** (2003), no. 1, 151–164.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT OSNABRÜCK, 49069 OSNABRÜCK, GERMANY
 Email address: dlevan@uos.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, 715 PATTERSON OFFICE TOWER,
 LEXINGTON, KY 40506-0027, USA
 Email address: uwe.nagel@uky.edu

INSTITUTE OF MATHEMATICS, VIETNAM ACADEMY OF SCIENCE AND TECHNOLOGY, 18 HOANG
 QUOC VIET, 10307 HANOI, VIETNAM
 Email address: ngdhop@gmail.com

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT OSNABRÜCK, 49069 OSNABRÜCK, GERMANY
 Email address: troemer@uos.de