

A NOTE ON THE NUMBER OF COMPLEX ROOTS OF A POLYNOMIAL OF ONE VARIABLE

CHU VAN TIEP¹

Abstract

The purpose of this article is twofold. The first aim is to find the number of complex solutions of a polynomial of one variable counting multiplicity. Then, our second aim is to approximate these solutions.

1 Introduction

From the Fundamental Theorem of Algebra, we deduce that every non-constant, single variable, degree n polynomial with real coefficients has counted with multiplicity, exactly n complex roots (see [3]). Mathematicians found the formula for the solution of a polynomial of degree less than 5 (see [2]). However, for an equation of degree greater than or equal to 5, we can't find a general formula of roots by radicals (see [4]). In this article, we want to find the number of complex solutions of a polynomial of one variable counting multiplicity. Then, we can approximate these solutions numerically. First of all, we list here some results about the information on the roots of a polynomial see for example [3, 6].

THEOREM 1.1 (DESCARTES' RULE [3]). The number of positive real roots of an equation with real coefficients is either equal to the number of its variations of sign or is less than that number by a positive even integer. A root of multiplicity m is here counted as m roots.

This rule can be derived as a corollary of the Budan's Theorem below.

THEOREM 1.2 (BUDAN'S THEOREM [3]). Let a and b be real numbers, $a < b$, neither a root of $f(x) = 0$, an equation of degree n with real coefficients. Let V_a denote the number of variations of the sign of

$$f(x); f'(x); f''(x); \dots; f^{(n)}(x)$$

for $x = a$, after vanishing terms have been deleted. Then $V_a - V_b$ is either the number of real roots of $f(x) = 0$ between a and b or exceeds the number of those roots by a positive even integer. A root of multiplicity m is here counted as m roots.

Given a square-free polynomial $f(x)$, denote:

- $f_0(x) = f(x)$.
- $f_1(x) = f'(x)$.
- $f_2(x) = -\text{rem}(f_0(x), f_1(x)) = f_1(x)q_0(x) - f_0(x)$, where $\text{rem}(f_0(x), f_1(x))$ is the remainder when polynomial long division is used to divide $f_0(x)$ by $f_1(x)$, and $q_0(x)$ is the quotient of said long division.

¹Faculty of Mathematics, The University of Danang - University of Science and Education, Da Nang 550000, Viet Nam
cvtiep@ued.udn.vn

- $f_3(x) = -\text{rem}(f_1(x), f_2(x)) = f_2(x)q_1(x) - f_1(x)$.
- In general, we define f_k inductively as $-\text{rem}(f_{k-2}(x), f_{k-1}(x)) = f_{k-1}(x)q_{k-2}(x) - f_{k-2}(x)$.
- Repeat this process until we arrive at some m such that $\text{rem}(f_{m-1}, f_m) = 0$, where $f_m(x) \neq 0$.

The finite sequence $f_0, f_1, \dots, f_m$ is called a Sturm sequence or a Sturm chain. Using this definition, we can locate the distinct real roots of a polynomial.

THEOREM 1.3 (STURM THEOREM [6]). Let $f(x) = 0$ be a polynomial equation and let

$$f_0(x), f_1(x), \dots, f_m(x)$$

be a Sturm sequence of the polynomial $f(x)$. Then for any two real numbers $a < b$, neither of which is a root of $f(x) = 0$, the number of distinct roots of $f(x) = 0$ lying between a and b equals the difference $g_f(a) - g_f(b)$ between the variations of the signs of the Sturm functions at $x = a$ and $x = b$.

2 Main result

In this section, we use the Sturm theorem to deduce a result on the number of complex solutions of a polynomial of one variable. For simplicity, for a polynomial $f(x)$ of degree n , we denote

- Rf : the number of real distinct roots of $f(x)$.
- $R^k f$: the number of real roots of multiple greater than or equal to k of $f(x)$.
- Cf : the number of complex distinct roots of $f(x)$.
- $C^k f$: the number of complex roots of multiple greater than or equal to k of $f(x)$.

We have the following theorem.

THEOREM 2.1. Consider $f(x) \in P_n[x]$ be a polynomial of degree $n \geq 1$. Let $f_0, f_1, \dots, f_m, f_{m+1} = 0$ be its Sturm sequence. Then, if f_m is a constant, then f only has simple roots. If f_m is a non-constant polynomial then f has multiple roots. Denote $f_{m_0}, f_{m_1}, \dots, f_{m_s}, f_{m_{s+1}}$ be the Sturm sequence of f_m . Then:

1. The number of real distinct roots of $f(x)$ is

$$R^1 f = Rf = g_f(-\infty) - g_f(+\infty) = g_{f_0/f_m}(-\infty) - g_{f_0/f_m}(+\infty).$$

2. The number of complex distinct roots of $f(x)$

$$C^1 f = Cf = \deg \frac{f_0}{f_m} - g(-\infty) + g(+\infty) = n - \deg f_m - Rf.$$

3. The number of real roots of $f(x)$ of multiplicity 1 is

$$g(-\infty) - g(+\infty) - g_{f_m}(-\infty) + g_{f_m}(+\infty) = Rf - Rf_m$$

where $g_{f_m}(x)$ the number of changes sign of Sturm sequence of $f_m(x)$.

4. The number of complex roots of multiple greater than or equal to 1 of $f(x)$ is

$$n - 2 \deg f_m + \deg f_{m_s} - R^1 f + R^1 f_m$$

where $f_{m_0}, f_{m_1}, \dots, f_{m_s}, f_{m_{(s+1)}} = 0$ is Sturm sequence of f_m .

Proof. 1. This is derived directly from the Sturm theorem since the multiplicity does not affect the difference $g_f(-\infty) - g_f(+\infty)$.

2. Since the polynomial f_0/f_m only contains simple roots of f so the numbers of distinct complex roots of f is

$$Cf = \deg \frac{f_0}{f_m} - g(-\infty) + g(+\infty).$$

3. If f_m (the greatest common divisor of f and its derivative) is a non-constant polynomial then $f(x)$ has roots with multiple greater than 1. We construct the Sturm sequence of f_m . Then $g_{f_m}(-\infty) + g_{f_m}(+\infty)$ is the numbers of distinct real roots of f_m . This is the number of roots with multiple greater than 1 of the $f(x)$. So, the number of real roots with multiplicity 1 is

$$g(-\infty) - g(+\infty) - g_{f_m}(-\infty) + g_{f_m}(+\infty) = Rf - Rf_m.$$

4. The number of distinct complex roots (not counting multiplicity) of f_m is

$$Cf_m = \deg \frac{f_m}{f_{m_s}} - g_{f_m}(-\infty) + g_{f_m}(+\infty) = \deg f_m - \deg f_{m_s} - Rf_m.$$

This is the number of complex roots of multiplicity greater than or equal 2 of $f(x)$. Thus, the the number of complex roots of multiplicity 1 of $f(x)$ is:

$$Cf - Cf_m = n - 2 \deg f_m + \deg f_{m_s} - R^1 f + R^1 f_m.$$

□

REMARK 2.2. Continue the Sturm sequence of f_{m_s} and so on, we will get the numbers of complex roots with higher multiplicity.

3 Numerical examples

In this section, we will use Newton's method² and the modified Newton's method to approximate the solution of $f(x)$. Newton's method is a root-finding algorithm that produces successively better approximations to the roots of a real-valued function. The most basic version starts with a real-valued function f , its derivative f' , and an initial guess x_0 for a root of f . If f satisfies certain assumptions and the initial guess is close, then

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

²also known as the Newton–Raphson method

is a better approximation of the root than x_0 . The process is repeated as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

until a sufficiently precise value is reached. If we know the multiplicity of roots, we can use the modified Newton's Method.

THEOREM 3.1 ([5]). If f is $(m+1)$ -times continuously differentiable on $[a, b]$, which contains a root r of multiplicity $m > 1$, then Modified Newton's Method

$$x_{i+1} = x_i - \frac{mf(x_i)}{f'(x_i)}$$

converges locally and quadratically to r .

REMARK 3.2. When $m = 1$, this theorem becomes the original Newton's Method.

Example 3.3. Consider the following polynomial of degree 8:

$$f(x) = x^8 + 12x^7 + 68x^6 + 236x^5 + 550x^4 + 884x^3 + 964x^2 + 660x + 225 = 0.$$

The Sturm sequence of f is:

$$f_0(x) = f(x) = x^8 + 12x^7 + 68x^6 + 236x^5 + 550x^4 + 884x^3 + 964x^2 + 660x + 225$$

$$f_1(x) = f'(x) = 8x^7 + 84x^6 + 408x^5 + 1180x^4 + 2200x^3 + 2652x^2 + 1928x + 660$$

$$f_2(x) = -5x^6 - 48x^5 - 215x^4 - 560x^3 - 903x^2 - 864x - 405$$

$$f_3(x) = x^5 + 5x^4 + 10x^3 + 6x^2 - 7x - 15$$

$$f_4(x) = x^4 + 6x^3 + 16x^2 + 22x + 15$$

$$f_5(x) = 0.$$

We have the following table:

x	$f_0(x)$	$f_1(x)$	$f_2(x)$	$f_3(x)$	$f_4(x)$	$g(x)$
$-\infty$	+	-	-	-	+	2
∞	+	+	-	+	+	2

So, the number of distinct real roots of $f(x)$ is

$$Rf = g(-\infty) - g(+\infty) = 2 - 2 = 0.$$

This means $f(x)$ has no real roots. So, all 8 roots are complex. We see that $f_4(x)$ is the greatest common divisor of f and f' . Thus, the number of distinct complex roots of $f(x)$ is

$$Cf = n - \deg f_4 - Rf = 8 - 4 - 0 = 4.$$

Next, we consider the sturm sequence of $f_4(x)$:

$$f_{4_0}(x) = f_4(x) = x^4 + 6x^3 + 16x^2 + 22x + 15$$

$$f_{4_1}(x) = 4x^3 + 18x^2 + 32x + 22$$

$$f_{4_2}(x) = -\frac{5}{4}x^2 - \frac{9}{2}x - \frac{27}{4}$$

$$f_{4_3}(x) = \frac{64}{25}x - \frac{64}{25}$$

$$f_{4_4}(x) = \frac{25}{2}$$

$$f_{4_5}(x) = 0.$$

We have

x	$f_{4_0}(x)$	$f_{4_1}(x)$	$f_{4_2}(x)$	$f_{4_3}(x)$	$f_{4_4}(x)$	$g(x)$
$-\infty$	+	-	-	-	+	2
∞	+	+	-	+	+	2

From the Sturm sequence of $f_4(x)$, we see that f_4 does not have multiple roots and from the table above we see that f_4 has no distinct real roots. Thus, f_4 have 4 distinct complex roots of the form: $a \pm ib$ và $c \pm id$ where $(a, b) \neq (c, d)$.


To approximate solutions of f , we use the modified Newton's method with $m = 2$. With initial guess $x_0 = 1 + i$, after 10 iterations, we get the result $x_{10} = -2 - i$. If we choose $x_0 = i$ after 7 iterations, we get the result : $x_7 = -1.0000 + 1.4142i$. If we use the Newton method, we need 17 iteration to get the same result as shown in the following tables:

k	Newton's method	Modified Newton's method
0	1+i	1+i
1	0.6210 + 0.9088i	0.2420 + 0.8175i
2	0.2766 + 0.8390i	-0.3888 + 0.7324i
3	-0.0403 + 0.7925i	-0.9568 + 0.7888i
4	-0.3364 + 0.7738i	-1.4765 + 1.1562i
5	-0.6163 + 0.7919i	-2.1792 - 0.7662i
6	-0.8792 + 0.8633i	-1.9572 - 0.9262i
7		-1.9989 - 1.0084i
8		-1.9999 - 1.0000i
9		-2.0000 - 1.0000i
10		-2.0000 - 1.0000i
17	-1.0000 + 1.4142i	
18	-1.0000 + 1.4142i	

and

k	Newton's method	Modified Newton's Method
0	i	i
1	-0.2800 + 0.9600i	-1.0334 + 1.0182i
2	-0.5371 + 0.9555i	-1.2977 + 1.4562i
3	-0.7684 + 0.9959i	-1.0302 + 1.2350i
4	-0.9588 + 1.0925i	-1.0413 + 1.4598i
5	-1.0662 + 1.2451i	-1.0048 + 1.4114i
6	-1.0419 + 1.3617i	-1.0000 + 1.4142i
7	-1.0190 + 1.3912i	-1.0000 + 1.4142i
8	-1.0091 + 1.4033i	
9	-1.0045 + 1.4089i	
10	-1.0022 + 1.4116i	
17	-1.0000 + 1.4142i	
18	-1.0000 + 1.4142i	

All 4 solutions are sketched in the following figure:



Example 3.4. Consider the polynomial equation of degree 11:

$$f(x) = x^{11} + x^{10} + x^9 + x^8 - 2x^7 - 2x^6 - 2x^5 - 2x^4 + x^3 + x^2 + x + 1 = 0,$$

The Sturm sequence of $f(x)$ is

$$f_0(x) = f(x) = x^{11} + x^{10} + x^9 + x^8 - 2x^7 - 2x^6 - 2x^5 - 2x^4 + x^3 + x^2 + x + 1$$


$$f_1(x) = p'(x) = 11x^{10} + 10x^9 + 9x^8 + 8x^7 - 14x^6 - 12x^5 - 10x^4 - 8x^3 + 3x^2 + 2x + 1$$

$$f_2(x) = -x^9 - 2x^8 + 8x^7 + 8x^6 + 10x^5 + 12x^4 - 8x^3 - 8x^2 - 9x - 10$$

$$f_3(x) = -x^8 + 2x^4 - 1$$

$$f_4(x) = -x^7 - x^6 - x^5 - x^4 + x^3 + x^2 + x + 1$$

$$f_5(x) = 0.$$



We have the following table:

x	$f_0(x)$	$f_1(x)$	$f_2(x)$	$f_3(x)$	$f_4(x)$	$g(x)$
$-\infty$	—	+	+	—	+	3
∞	+	+	—	—	—	1

Thus, f has $3 - 1 = 2$ distinct real solution (not counting multiplicity). The number of complex distinct roots of f is

$$11 - 7 - 2 = 2.$$

So f has two complex roots (not counting multiplicity) of the form $a \pm ib$. To get the information about multiplicity, we consider the Sturm sequence of f_4 : $f_4(x)$:

$$f_{40}(x) = f_4(x) = -x^7 - x^6 - x^5 - x^4 + x^3 + x^2 + x + 1$$

$$f_{41}(x) = f_4'(x) = -7x^6 - 6x^5 - 5x^4 - 4x^3 + 3x^2 + 2x + 1$$

$$f_{42}(x) = x^5 + 2x^4 - 4x^3 - 4x^2 - 5x - 6$$

$$f_{43}(x) = x^4 - 1$$

$$f_{44}(x) = x^3 + x^2 + x + 1$$

$$f_{45}(x) = 0.$$

So, we have the following table:

x	$f_{40}(x)$	$f_{41}(x)$	$f_{42}(x)$	$f_{43}(x)$	$f_{44}(x)$	$g(x)$
$-\infty$	+	—	—	+	0	2
∞	+	+	+	+	0	0

Thus, $f_4(x)$ has $2 - 0 = 2$ distinct real roots (not counting multiplicity). Then, we deduce that the roots f have multiplicity greater than or equal to 2. Then, f has no simple real root and f two complex roots with multiplicity greater than or equal to 2. We can continue to calculate the Sturm sequence of f_{43} . However, in this particular case, we have $f_{43}(x) = (x^2 + 1)(x + 1)$. Thus,

it has one simple root $x = -1$ and two complex roots $x = \pm i$. Therefore, $x = -1$ is a root with multiplicity 3, and $x = \pm i$ are also roots with multiplicity 3. The other real root is multiple 2. Then $f(x) = (x + i)^3(x - i)^3(x + 1)^3(x - x_0)^2$. Divide $f(x)$ by $(x + i)^3(x - i)^3(x + 1)^3 = (x + 1)^3(x + 1)^3$ we derive $x_0 = 1$. So, we have all the roots

- $x = 1$ (multiple 2)
- $x = -1$ (multiple 3)
- $x = \pm i$ (multiple 3)

In the general case, when it is not easy to factor $f_{4_3}(x)$, we continue this process to get the result.

4 Conclusion

The main result of the article is to state the theorem for calculating the number of complex solutions with corresponding multiples of any n degree algebraic equation. Then, we can accurately calculate the solution in special cases or approximate the solution with Newton's algorithm with known multiples.

Acknowledgement. Part of this work was done while the author was visiting the Vietnam Institute for Advanced Study in Mathematics (VIASM). The author would like to thank the VIASM for financial support and hospitality.

References

- [1] Ackleh, A.S. and Allen, E.J. and Kearfott, R.B. and Seshaiyer, P. (2009), *Classical and Modern Numerical Analysis: Theory, Methods and Practice*, CRC Press.
- [2] Cardano, G. and Witmer, T.R. (1993), *Ars Magna Or The Rules of Algebra*, Dover.
- [3] Leonard E. Dickson (2009), *First course in the theory of equations*, University of Michigan Library.
- [4] Jean-Pierre Tignol (2016), *Galois' Theory of Algebraic Equations*, World Scientific Publishing Company.
- [5] Richard L Burden; J Douglas Faires (2011), *Numerical analysis*, Brooks/Cole Cengage Learning.
- [6] K.T.Leung, I.A.C Mok, S.N.Suen (1993), *Polynomials and Equations*, Hong Kong University Press.
- [7] Sturm, Jacques Charles François (1829), *Mémoire sur la résolution des équations numériques*, Bulletin des Sciences de Férussac.